

Motor Function Recovery System for Stroke Patients

Published date: Sept. 11, 2018

Technology description

Background: A stroke occurs when blood flow to an area in the brain is cut off. Brain cells deprived of oxygen and glucose die and, in many cases, permanent brain damage can result. Strokes are the leading cause of serious, long-term disability in the United States. The ultimate goal of rehabilitation is to improve motor function so that the stroke survivor can become as independent as possible. This must be accomplished in a way that preserves dignity and motivates the survivor to relearn basic skills that the stroke may have impaired - skills like bathing, eating, dressing and walking.

Technology: Georgia Tech researchers have developed a method to stimulate certain areas of the body to improve motor function post-stroke. The stimulation is delivered through an apparatus that can be worn or placed against the body (such as an arm rest). The vibration is non-focal, meaning it is not specific to a part of the body or muscle. Vibration motors are sewn into the fabric of a wearable device or mounted on the surface of a stationary device. The patient can either wear the device on their body or rest in contact with the tactile stimulation device—no further exercise is needed. Several specially engineered stimulation patterns provide even coverage. This device and method aid in the transition from hospital to full recovery, acting as an alternative to traditional methods of treatment.

Application area

Stroke recovery

Mobile rehab devices

Advantages

Unconventional – Passive stimulation unlike traditional therapy or rehabilitation

Tactile – No electrostimulation, more comfortable to patients

Convenient – Little to no effort/attention required

Cheaper – Mobile rehab outside of a clinic or hospital

Institution

Georgia Institute of Technology

Caitlyn Seim

联系我们



叶先生

电话: 021-65679356 手机: 13414935137

邮箱: yeyingsheng@zf-ym.com