

Campylobacter detection and strain identification

Published date: Aug. 12, 2009

Technology description

Development of a technology for a simple, sensitive, Campylobacter detection and strain identification kit that generates specific, reproducible results.

Description

Researchers affiliated with the National Microbiology Laboratory (NML) in Winnipeg, Manitoba, Canada have developed the technology for a simple, sensitive, Campylobacter detection and strain identification kit that generates specific, reproducible results.

The Challenge

Campylobacter is the leading cause of bacterial gastroenteritis in the developed world.

Symptoms of Campylobacter infection are diarrhea, abdominal cramps, vomiting, and fever. Dysentery may also occur.

Campylobacter can be found in food and water. Poultry is the major source of human Campylobacter infection. Between 2.1 and 2.4 million cases of human illness caused by the Campylobacter microorganism occur in the United States yearly, and the death toll from the pathogen approaches 1,000.

Detection of the Campylobacter pathogen is crucial to preventing Campylobacter spread, treating Campylobacter disease, and establishing the epidemiology of Campylobacter infection. Although the disease is sporadic in distribution, outbreaks have been reported.

Identification of Campylobacter strains is important for tracing the transmission history of the strain or strains causing disease and for facilitating the recall of contaminated food.

Most Campylobacter detection and strain identification technologies are less than ideal. Campylobacter is difficult to grow, necessitating detection technology not overly dependant on sample amplification. Detection sensitivity is an issue. Those technologies that can identify strains employ Restriction

Fragment Length Polymorphism (RFLP) or DNA probe techniques, and because these techniques are complex, widespread use of presently available strain identification technologies is problematic.

An easy-to-use, sensitive, rapid *Campylobacter* detection kit that simultaneously detects and identifies all species of *Campylobacter* with a high degree of sensitivity is urgently needed.

Application area

The NML kit's ease of use, sensitivity, and reliability makes widespread use of it at many levels of medical practice, medical research, and food and water inspection possible and practical. It has civilian and military applications and, potentially, a worldwide market.

Advantages

The kit can, in three hours, simultaneously detect *Campylobacter* and identify the strains of *Campylobacter* most often responsible for *Campylobacter* disease even when the pathogens are present at very low levels.

It does not use the complex RFLP or DNA probe techniques.

No other *Campylobacter* detection technology simultaneously detects and identifies so many species of *Campylobacter*.

The NML kit uses a Multiplex Polymerase Chain Reaction (PCR) assays to simultaneously detect in a single assay genes that identify all strains of the *Campylobacter* species *C. jejuni*, *C. coli*, *C. lari*, *C. upsaliensis*, and *C. fetus*.

A double-stranded DNA helix can be separated longitudinally, and each strand that results from the split can serve as a template for growth of a DNA strand identical to the one from which the template.

The kit contains primers, each designed to attach to a separated DNA strand. If a DNA strand from one of the targeted genes is present in the sample being tested, the appropriate primer will attach to it and a DNA strand will grow. This growth indicates the presence of the gene and the *Campylobacter* strain associated with it.

The NML kit targets the genes *hipO* and 23S rRNA to detect *C. jejuni*, the gene *glyA* to detect *C. coli*, *C. lari*, and *C. upsaliensis*, and the gene *sapB2* to detect *C. fetus*.

The kit contains an amplification mixture that facilitates DNA strand growth and a positive control that confirms the test conditions are conducive to such growth.

Samples, such as food or stool specimens, suspected of contamination with any or all of the *Campylobacter* species known to cause disease may be tested directly in a single assay and do not require preparation or pathogen isolation.

Institution

[Public Health Agency of Canada \(PHAC\) - Agence de la santé publique du Canada \(ASPC\)](#)

联系我们



叶先生

电话 : 021-65679356

手机 : 13414935137

邮箱 : yeyingsheng@zf-ym.com