

Measuring the Activity of a Specific Fraction of Albumin for Early Diagnosis of COPD or Sepsis

Published date: Feb. 1, 2012

Technology description

Description

Chronic obstructive pulmonary disease (COPD) is a complex group of conditions usually related to cigarette smoking. COPD is associated with progressive airway obstruction and loss of lung function. This disease is the fourth leading cause of death in the U.S., and caring for patients with COPD costs as much as 40 billion dollars each year.

Current methods of diagnosing COPD include pulmonary function testing, pulse oximetry, radiological procedures and arterial blood gases. However, these tests only detect relatively advanced disease. Early diagnosis is critical to slowing disease progression and preventing irreversible lung damage.

The level of inflammation in patients with COPD may provide a means of early diagnosis. However, a recent study of 36 systemic biomarkers found that they were not useful in predicting which patients will develop severe respiratory dysfunction. Currently, no reliable way exists to predict which individuals who smoke tobacco will develop COPD or which patients with COPD will become progressively worse.

Sepsis is a life-threatening systemic inflammatory response syndrome. Patients with the onset of an infection are at risk of developing sepsis. Like COPD, early diagnosis is critical to improving patient outcomes, but accurate ways of detecting sepsis at an early stage currently are not available.

This assay measures the functional activity of a specific fraction of albumin (SFA) in serum or plasma. The assay uses a liposome that contains a fluorescently labeled fatty acid and a negatively charged phospholipid as a substrate. The liposome is mixed with phospholipase A₂ (PLA₂), an enzyme that plays a key role in inflammation, and a biological sample from a patient. The SFA in the sample then acts to remove the labeled fatty acid from the liposome, causing a detectable change in fluorescence intensity. This change can be used to determine SFA activity.

The measured SFA activity then can be compared to a normal range of SFA activity or to a baseline measurement from the patient to determine if it is decreasing, abnormally low or normal. A decrease or low measurement of SFA activity can indicate inflammation. If the patient is a tobacco smoker, a decrease in activity as compared to an earlier measurement from the patient suggests the patient has developed or will soon develop COPD. If a patient has recently undergone surgery, a decrease in SFA activity can indicate infection and the possibility of sepsis. On the other hand, a return of SFA activity to normal levels may indicate that the patient has recovered.

Advantages

UW-Madison researchers have developed a sensitive and rapid high throughput assay that can be used for early detection of COPD or sepsis.

Institution

[University of Wisconsin](#)

联系我们



叶先生

电话 : 021-65679356

手机 : 13414935137

邮箱 : yeyingsheng@zf-ym.com